

A precise determination of the decay constant of the D_s -meson in quenched QCD



Andreas Jüttner^a and Juri Rolf^b

Institut für Physik, Humboldt-Universität zu Berlin

Invalidenstr. 110, D-10115 Berlin, Germany

Email: ^ajuettner@physik.hu-berlin.de

^brolf@physik.hu-berlin.de

Abstract

We present a lattice determination of the leptonic decay constant F_{D_s} of the D_s -meson using its mass and F_K as experimental input. Setting the scale with $F_K = 160$ MeV, our final result is $F_{D_s} = 252(9)$ MeV. The error contains all uncertainties apart from the quenched approximation. Setting the scale with the nucleon mass instead leads to an decrease of $20(1)$ MeV of the leptonic decay constant.

Keywords: lattice QCD, leptonic decay constant, heavy-light mesons, CKM-matrix

PACS: 11.15.Ha, 12.38.Gc, 13.20.Fc, 14.40.Lb

1 Introduction

The D_s^+ -meson consists of a c and a $\bar{s}$ quark. It is stable in QCD and decays by an emission of a W^+ -boson into a lepton and a neutrino. The amplitude of this decay is characterized by the decay constant F_{D_s} which is defined by the QCD matrix element

$$\langle 0 | A_\mu(0) | D_s(p) \rangle = i p_\mu F_{D_s} \quad (1.1)$$

of the axial current $A_\mu = \bar{s} \gamma_\mu \gamma_5 c$.

Given the CKM-matrix element V_{cs} , F_{D_s} can be measured experimentally by studying the branching ratio $\text{BR}(D_s \rightarrow l \nu)$. The current status of the experimental determination of F_{D_s} has been summarized in [1]. Currently, the most precise data come from the ALEPH experiment [2], $F_{D_s} = (285 \pm 19 \pm 40) \text{ MeV}$, and from CLEO [3], $F_{D_s} = (280 \pm 17 \pm 25 \pm 34) \text{ MeV}$. In the next couple of years CLEO aims at reducing their error to two percent precision [4].

The status of lattice computations for F_{D_s} [5,6,7] was reviewed in [8,9] with a world average of $F_{D_s} = 255 \pm 30 \text{ MeV}$. With QCD sum rules one currently gets $F_{D_s} = 235 \pm 24 \text{ MeV}$ [10,11].

The goal of this work is a computation on the lattice of the weak decay constant F_{D_s} to 3 percent accuracy up to the quenched approximation. All the systematic and statistical errors will be analyzed and we will also estimate the size of the quenched scale ambiguity. A precise quenched calculation of F_{D_s} together with precise experimental data supplies us with a test of quenched lattice QCD. This is of importance for the determination of the B-meson decay constant f_B from lattice QCD which is an essential input in the measurement of the third generation CKM-matrix elements.

2 Strategy

The decay constant F_{D_s} is defined in terms of a QCD matrix element. To evaluate this matrix element we have to eliminate the bare parameters of the QCD Lagrangian, that is the bare gauge coupling g_0 and the bare masses of the relevant quarks, in favour of physical observables. One possible hadronic scheme, which has been summarized in [12] and developed in [13,14,12] is to use the decay constant $F_K = 160 \text{ MeV}$ to set the scale¹ and eliminate the bare masses of the strange and charm quarks in favour of the masses of the K and the D_s -meson. Neglecting isospin breaking, the quark mass ratio $M_s/\hat{M} = 24.4 \pm 1.5$ with $\hat{M} = \frac{1}{2}(M_u + M_d)$ is taken from chiral perturbation theory [15].

The results in [13,14,12] are given in terms of the Sommer scale r_0 which is derived from the force between static colour sources. r_0/a has been computed as a function of the bare coupling g_0 to a high precision for a wide range of cutoff values [16,17,18]. It is only affected

¹i.e. to compute the lattice spacing a in physical units as a function of the bare coupling

by lattice artifacts of $O(a^2)$ in the quenched theory [18]. The relation to other hadronic scales is known [14], $r_0 F_K = 0.415(9)$, and $r_0 m_N \approx 2.6$. This illustrates the inconsistency of the quenched approximation since the ratio F_K/m_N deviates by approximately 10% from its experimental value. This is the typical size of the quenched scale ambiguity. Setting the scale with $F_K = 160$ MeV corresponds to setting $r_0 = 0.5$ fm while $m_N = 938$ MeV corresponds to $r_0 = 0.55$ fm. Below we will estimate the size of the quenched scale ambiguity for the weak decay constant of the D_s -meson.

This work aims at a scaling study of F_{D_s} on the lattice. Thus we keep the physical conditions constant and only vary the lattice spacing. We perform numerical simulations in $O(a)$ improved lattice QCD using Schrödinger functional boundary conditions [19,20] on a $L^3 \times T$ space time cylinder. The practical advantages of this approach are discussed in [21]. For unexplained notation we refer to [22].

In particular we define the meson sources

$$\mathcal{O} = a^6 \sum_{\mathbf{y}, \mathbf{z}} \bar{\zeta}_j(\mathbf{y}) \gamma_5 \zeta_i(\mathbf{z}), \quad \mathcal{O}' = a^6 \sum_{\mathbf{y}, \mathbf{z}} \bar{\zeta}'_j(\mathbf{y}) \gamma_5 \zeta'_i(\mathbf{z}), \quad (2.2)$$

with flavour indices $i \neq j$ at the $x_0 = 0$ and the $x_0 = T$ boundary time slices, respectively. The correlation functions

$$f_A(x_0) = -\frac{1}{2} \langle \mathcal{O} A_0(x) \rangle, \quad f_P(x_0) = -\frac{1}{2} \langle \mathcal{O} P(x) \rangle, \quad f_1 = -\frac{1}{3L^6} \langle \mathcal{O}' \mathcal{O} \rangle \quad (2.3)$$

are then used to compute the decay constant F_{D_s} at finite lattice spacing. Here $A_\mu(x) = \bar{\psi}_i(x) \gamma_\mu \gamma_5 \psi_j(x)$ is the axial current and $P(x) = \bar{\psi}_i(x) \gamma_5 \psi_j(x)$ is the pseudoscalar density. The leading $O(a)$ cutoff effects in f_A are canceled by using a clover improved action [23] with coefficient c_{sw} and by improving the axial current,

$$A_\mu^I(x) = A_\mu(x) + c_A a \tilde{\partial}_\mu P(x). \quad (2.4)$$

Both coefficients c_{sw} and c_A are known non perturbatively in the quenched approximation for bare couplings $g_0^2 = 6/\beta \leq 1$ [24]. The improved correlation function f_A^I is then defined in analogy to f_A .

On the lattice the axial current receives a multiplicative renormalization by the finite factor Z_A . The renormalized axial current is thus given by

$$(A_R)_\mu = Z_A (1 + b_A (am_{q,i} + am_{q,j})/2) A_\mu^I + O(a^2), \quad (2.5)$$

where the $O(a)$ artifacts that are proportional to the bare subtracted quark masses am_q are canceled by the term proportional to b_A .

The correlation functions f_A^I and f_P decay exponentially proportional to $e^{-x_0 m_{D_s}}$ when the excitations due to higher states are small enough. Thus the meson mass m_{D_s} , which has the experimental value $r_0 m_{D_s} = 4.988$ [25], can be obtained from the plateau average of the

effective pseudoscalar mass

$$am_{\text{PS}}(x_0 + a/2) = \log \frac{f_A^I(x_0)}{f_A^I(x_0 + a)}. \quad (2.6)$$

The weak decay constant of the D_s -meson at finite lattice spacing can then be obtained through ²

$$F_{D_s} = -2Z_A(1 + b_A(am_{q,i} + am_{q,j})/2) \frac{f_A^I}{\sqrt{f_1}} (m_{D_s}L^3)^{-1/2} e^{(x_0 - T/2)m_{D_s}} \\ \times \{1 - \eta_A^{D_s} e^{-x_0\Delta} - \eta_A^0 e^{-(T-x_0)m_G}\} + O(a^2). \quad (2.7)$$

Here the factor $(m_{D_s}L^3)^{-1/2}$ takes into account the normalization of one particle states. The contribution $f_1^{-1/2}$ cancels out the dependence on the meson sources. Because of the exponential decay of the correlation function f_A^I the product in (2.7) is expected to exhibit a plateau at intermediate times when the contribution $\eta_A^{D_s} e^{-x_0\Delta}$ of the first excited state and the contribution $\eta_A^0 e^{-(T-x_0)m_G}$ from the O^{++} glueball both are small. A plateau average can then be performed to increase the signal and is understood in (2.7). Further explanations for equation (2.7) and details can be found in [21].

3 Numerical results

3.1 Parameters

Our choice of simulation parameters is shown in table 1. The bare couplings and the hopping

β	n_{meas}	L/a	L/r_0	κ_{critical}	κ_s	κ_c	$r_0 m_{D_s}$
6.0	380	16	2.98	0.135196	0.133929	0.119053	4.972(22)
6.1	301	24	3.79	0.135496	0.134439	0.122490	4.981(23)
6.2	251	24	3.26	0.135795	0.134832	0.124637	5.000(25)
6.45	289	32	3.06	0.135701	0.135124	0.128131	5.042(29)

Table 1: Statistics and parameters for our simulations and demonstration of constant physical conditions.

parameters for the strange quark are the same as used in [12], and the hopping parameters for the charm quark have been found in that work. The non perturbatively defined improvement coefficients c_{sw} and c_A are taken from [24]. b_A has been obtained non perturbatively in [26]. Since it is difficult to compute (cmp. [27]) and since we have to extrapolate their results slightly we have used one-loop perturbation theory [28] as well. The conversion to physical

²In [21] the first minus sign is missing.

units is done with the Sommer scale r_0 . From the fit function of [16] we infer that our lattice size $L/a = 16$ at $\beta = 6.0$ corresponds to $L/r_0 = 2.98$. The results of [14] show that at this volume the finite size effects may be neglected. The lattice spacing a is varied from 0.1 fm to 0.05 fm.

Furthermore we choose $T/L = 2$ in contrast to $T/L = 2.5$ used for the computation of the charm quark mass in [12] to reduce rounding effects, which would be the dominant systematic effects on F_{D_s} otherwise.

The fourth column of table 1 shows that the above choice of simulation parameters corresponds to approximately constant physical box sizes. Also the mass of the D_s -meson takes its correct value up to small fluctuations. We have checked that these lead to negligible corrections for the decay constant F_{D_s} .

3.2 Computation of the D_s -meson decay constant

To compute the meson mass m_{D_s} and the decay constant F_{D_s} we calculate the combinations (2.6,2.7) of correlation functions. For all the parameter choices we find plateaus as functions of x_0 . Our task is to find the plateau region such that the effect of excited states is negligible compared to our statistical error.

When dealing with heavy quark propagators on single precision machines there is always the danger of roundoff problems which might be the dominant uncontrolled systematic error. Since the computation of the decay constant F_{D_s} involves propagators extending from the $x_0 = 0$ to the $x_0 = T$ boundary in the correlation functions f_1 the discussion of these issues becomes even more important as for the computation of masses. From [12] we estimate that for $T = 2L$ instead of $T = 2.5L$ taken there the rounding errors for F_{D_s} should be small enough. To show this we computed our observables on $\mathcal{O}(10)$ configurations at $\beta = 6.0$ and 6.45 in single as well as in double precision. This check reveals that the rounding errors for F_{D_s} are of the order of one per mill.

From fits of $F_{D_s}(x_0)$ using an a priori chosen plateau range we extract the contributions $\eta_A^{D_s} e^{-x_0 \Delta}$ and $\eta_A^0 e^{-(T-x_0)m_G}$ of the first excited state and the 0^{++} glueball excitation, respectively. The theory in [21] predicts relations between the prefactors and the corresponding prefactors for the pseudoscalar masses which are roughly ³ fulfilled.

All the systematic effects that deteriorate the plateau are added. With an a priori chosen upper bound δ for the maximal relative systematic error in a point $F_{D_s}(x_0)$ we define the time interval for the plateau region such that for all times in this interval the added systematic contribution is smaller than δ . The systematic error for the plateau averaged observable is then even smaller. For the decay constant we took $\delta = 0.5\%$. The plateaus defined in this way extend from $4r_0$ to $5r_0$ at the four β values considered. For the pseudoscalar mass we used $\delta = 0.3\%$.

³In some cases the fit function is not determined precisely. Then we choose the fit function such that the contribution of the excited states is rather overestimated. Note that we do not attempt to measure Δ or $\eta_A^{D_s}$ but rather want to estimate our systematic effects.

3.3 Continuum limit

We perform the plateau averages of (2.7) and obtain the decay constant at finite lattice spacings as shown in table 2. These data can be extrapolated to the continuum limit,

β	$aF_{D_s}^{\text{bare}} _{c_A=0}$	$\partial aF_{D_s}^{\text{bare}}/\partial c_A$	$r_0F_{D_s}$
6.0	0.1144(25)	0.2375(58)	0.540(14)
6.1	0.0979(17)	0.1699(33)	0.576(13)
6.2	0.0863(20)	0.1280(32)	0.598(16)
6.45	0.0633(14)	0.0664(17)	0.614(15)

Table 2: Simulation results for F_{D_s} .

leading to our main result. Since we employ $O(a)$ improvement, the natural ansatz for the continuum extrapolation is linear in a^2 . This extrapolation is shown in figure 1. In the fit




Figure 1: Continuum extrapolation of F_{D_s} .

we neglect the point at $\beta = 6.0$ which is farthest away from the continuum limit. We quote

$$r_0F_{D_s} = 0.638(24) \quad (3.8)$$

as our final result. Using $r_0 = 0.5 \text{ fm}$ this corresponds to $F_{D_s} = 252(9) \text{ MeV}$. Using 1-loop perturbation theory for b_A instead of the results of [26] leads to

$$r_0F_{D_s} = 0.631(24), \quad (3.9)$$

corresponding to $F_{D_s} = 249(9)$ MeV.

3.4 Quenched scale ambiguity

To estimate the quenched scale ambiguity we consider $r_0 F_{D_s}$ as a function $f(z)$ of $z = r_0 m_{D_s}$ around the physical value of z . This is possible since in addition to the hopping parameters leading to the physical strange and charm quark masses we have computed our observables at five more hopping parameter values around the hopping parameter of the charm quark for each bare coupling considered. Here we will only use the two hopping parameters closest to the charm value and present further results in a separate publication.

We expand $f(z) = f(z_0) + (z - z_0)f'(z_0) + \dots$ in a Taylor series around $z_0 = 4.988$. $f'(z_0)$ is estimated from a linear fit of $r_0 F_{D_s}$ as a function of $r_0 m_{D_s}$. A 10% increase of r_0 corresponds to $z - z_0 = 0.5$. Our estimate of the corresponding effect on $r_0 F_{D_s}$ in the continuum limit is

$$0.5f'(z_0) = 0.008(3). \quad (3.10)$$

Converting back to physical units (now using $r_0 = 0.55$ fm) we estimate that F_{D_s} decreases by 20 MeV corresponding to eight percent.

4 Conclusions

Our main result is $r_0 F_{D_s} = 0.638(24)$. Using $r_0 = 0.5$ fm this corresponds to $F_{D_s} = 252(9)$ MeV. This is compatible with all the experimental data currently available and with most theoretical estimates as well. The precision of our result however matches the goal of precision of future experiments, for example CLEO.

Under a scale shift of 10%, which is typical for quenched spectrum computations F_{D_s} decreases by eight percent. Assuming that all the scale ambiguity comes from neglecting dynamical sea quark effects this might also give an idea of the unquenching effect for F_{D_s} which however has to be probed in an unquenched simulation.

Since our complete data set consists also of simulations at larger quark masses we will be able to look at the functional dependence of the weak decay constant around the charm quark mass. Together with a new computation of the decay constant in the static approximation we will then interpolate to the B -meson mass and compute F_B with a better precision than is currently available. The corresponding analysis has still to be done and further simulations will be necessary so that these exciting results will be presented in another publication.

This work is part of the ALPHA collaboration research programme. It was supported by the European Community under the grant HPRN-CT-2000-00145 Hadrons/Lattice QCD and by the Deutsche Forschungsgemeinschaft in the SFB/TR 09 and the Graduiertenkolleg GK271. All the production runs were carried out on machines of the APE1000 series at DESY-Zeuthen. The checks of the rounding errors in double precision were done with a C-

code based on the MILC collaboration's public lattice gauge theory code, see [29]. This code ran on the IBM p690 system of the HLRN [30] and on the PC cluster at DESY-Zeuthen. We thank the staff at the computer centres for their help, and M. Della Morte, S. Sint, R. Sommer, and U. Wolff for useful discussions.

References

- [1] S. Söldner-Rembold, JHEP Proceedings, HEP2001, hep-ex/0109023.
- [2] A. Heister et. al., Phys. Lett. B **528** (2002) 1.
- [3] M. Chadha et. al., Phys. Rev. D58:032002 (1998).
- [4] Project description for CLEO-c,
<http://www.lns.cornell.edu/public/CLEO/spoke/CLEOc/ProjDesc.html>.
- [5] C. W. Bernard *et al.*, Nucl. Phys. Proc. Suppl. **94** (2001) 346.
- [6] D. Becirevic, Nucl. Phys. Proc. Suppl. **94** (2001) 337.
- [7] C. M. Maynard [UKQCD Collaboration], Nucl. Phys. Proc. Suppl. **106** (2002) 388.
- [8] T. Draper, Nucl. Phys. Proc. Suppl. **73** (1999) 43.
- [9] C. W. Bernard, Nucl. Phys. Proc. Suppl. **94** (2001) 159.
- [10] S. Narison, Phys. Lett. B **520** (2001) 115.
- [11] A. A. Penin and M. Steinhauser, Phys. Rev. D **65** (2002) 054006.
- [12] J. Rolf and S. Sint [ALPHA Collaboration], JHEP 12 (2002) 007.
- [13] S. Capitani, M. Lüscher, R. Sommer and H. Wittig [ALPHA Collaboration], Nucl. Phys. B **544** (1999) 669.
- [14] J. Garden, J. Heitger, R. Sommer and H. Wittig [ALPHA Collaboration], Nucl. Phys. B **571** (2000) 237.
- [15] H. Leutwyler, Phys. Lett. B **378** (1996) 313.
- [16] M. Guagnelli, R. Sommer and H. Wittig, Nucl. Phys. B **535** (1998) 389.
- [17] R. G. Edwards, U. M. Heller and T. R. Klassen, Nucl. Phys. B **517** (1998) 377.
- [18] S. Necco and R. Sommer, Nucl. Phys. B **622** (2002) 328.
- [19] M. Lüscher, R. Narayanan, P. Weisz and U. Wolff, Nucl. Phys. B **384** (1992) 168.

- [20] S. Sint, Nucl. Phys. B **421** (1994) 135.
- [21] M. Guagnelli, J. Heitger, R. Sommer and H. Wittig [ALPHA Collaboration], Nucl. Phys. B **560** (1999) 465.
- [22] M. Lüscher, S. Sint, R. Sommer and P. Weisz, Nucl. Phys. B **478** (1996) 365.
- [23] B. Sheikholeslami and R. Wohlert, Nucl. Phys. B **259** (1985) 572.
- [24] M. Lüscher, S. Sint, R. Sommer, P. Weisz and U. Wolff, Nucl. Phys. B **491** (1997) 323.
- [25] K. Hagiwara et al. [Particle Data Group Collaboration], Phys. Rev. D **66** (2002) 010001.
- [26] T. Bhattacharya, R. Gupta, W. j. Lee and S. Sharpe, Nucl. Phys. Proc. Suppl. **106** (2002) 789.
- [27] M. Guagnelli, R. Petronzio, J. Rolf, S. Sint, R. Sommer and U. Wolff [ALPHA Collaboration], Nucl. Phys. B **595** (2001) 44.
- [28] S. Sint and P. Weisz, Nucl. Phys. B **502** (1997) 251.
- [29] <http://www.physics.indiana.edu/~sg/milc.html>.
- [30] <http://www.hlrn.de>.